

Innovative Optimal Approach to Implement Fast Propagation Tissue Culture

Hossein Nazarian¹ and Amir Hossein Ghadimi^{2*}

¹Department of Agricultural Biotechnology, Payame Noor University, Karaj, Iran.

²Department of Food Science and Technology, Yasooj Branch, Islamic Azad University, Iran.

<http://dx.doi.org/10.13005/bbra/2485>

(Received: 06 April 2017; accepted: 06 May 2017)

In this study bulbetets of royal lily have been used as explants. The innovative approach is based on optimal placement and adventitious sprout induction and the adventitious sprout proliferation in special conditions. Then effects of different parts of scales and placement method, types and concentrations of cytokinin, different parts of bulbetet scales and placement method on adventitious sprout induction have been discussed. At the end the effects of plant growth regulators on induction and multiplication of bulbetet scales of royal lily have been studied. The results represents that the regeneration ability of outer layer scales of royal lily is better than the middle ones, which was better than the inner ones. The adaxial side of royal lily bulbetets in contact with medium upward can produce more and stronger adventitious sprouts than the adaxial side downward.

Keywords: Optimal placement, tissue culture, fast propagation, adventitious sprout.

Royal lily, flowering bulb, is one of the most important species of Liliaceae and *Lilium*, which was discovered in Mingjiang region, Sichuan province by famous botanist Wilson EH in 1903 and was introduced to Britain and America in 1905 and then became the worldwide cultivated variety¹⁹.

In production, the common propagation method of Lily is division propagation. But the reproductive rate of division propagation is not only too low but also, the seedball is prone to degeneration due to accumulative diseases, giving rise to an adverse effect on growth and ornamental values²⁰. Although, traditional asexual propagation like scale cutting is available for some lily varieties it is not frequently used because the scales are

easily rotten after cutting [13]. Using tissue culture method can get not only high propagation coefficient, but also virus elimination and it can be also combined with genetic engineering and provide a new method for plant breeding⁹.

Many researchers have studied culture in vitro of lilies⁷. Cultivated varieties were studied more and wild species were less focused on²¹, especially Royal lily. Currently, there are only two cases of reports on culture in vitro of Royal lily seeds¹⁴ and bulbs¹⁸. In most of previous researches, bulbs^{12,8}, leaves^{1,17} and roots⁵ have been used as explants, but the explants are limited to get and meanwhile a higher contamination rate is difficult to overcome. The bulbils are on the plant aerial parts, carrying less amount of germs, not easy to be damaged, seldom develop a phenomenon of endogenous bacterial in subculture³. Moreover, bulbils of Royal lily are enough in quantity, plump and easy to get. Picking bulbils does not affect not

* To whom all correspondence should be addressed.
E-mail: ahghadimi67@gmail.com

only on the development of bulbs and flower bud differentiation, but also plant growth and flowering. Therefore, there are many advantages in using bulbils as explants.

Using bulbils as explants, the aim of the present study was to establish a more perfect in vitro regeneration system and provide theoretical and technical basis for fast propagation, detoxification utilization and resource conservation of Royal lily.

MATERIALS AND METHODS

The bulbils of Royal lily were obtained from Iran, Mahallat. The experiments were carried out in the Comprehensive Laboratory of agricultural and biotechnology in Payame Noor university of Iran.

Adventitious bud induction

The bulbils of Royal lily were rinsed with a little detergent and two drops Twain 20 and washed in running water for 1 h. Then they were surface sterilized with 75% alcohol for 20 sec first, rinsed twice in sterile distilled water and then sterilized with 0.1% HgCl₂ for 12 min followed by three final rinses in sterile distilled water. The bulbils were divided into three parts: the outer layer (outer 1~2 layers), the middle layer (outer 3~4 layers) and the inner layer (the rest). The scales were latitudinally cut for three times and inoculated by means of adaxial side upward and adaxial side downward. The medium was MS supplemented with NAA 0.2 mg L⁻¹, Casein Hydrolysate (CH) 500 mg L⁻¹, Sucrose 20 g L⁻¹, Agar 9 g L⁻¹ and different concentrations of BA, KT and ZT, respectively. The PH was adjusted to 5.8. Data on explants forming adventitious bud were observed after 35 days.

Adventitious bud proliferation

Adventitious buds excised from in vitro scales were subcultured on MS medium containing BA (0.5, 1.0 and 2.0 mg L⁻¹), KT (1.0 and 1.5 mg L⁻¹) and NAA (0.1 and 0.2 mg L⁻¹). All the media contained casein hydrolysate (CH) 500 mg L⁻¹, Sucrose 20 g L⁻¹, Agar 9 g L⁻¹ and were adjusted to pH 5.8. After 6 weeks of culture, the number and height of newly formed adventitious buds were recorded.

Culture conditions

In addition to the special statement,

cultures were incubated at 25°C under 14 h photoperiod provided by white fluorescent lights.

RESULTS AND DISCUSSION

Effects of different parts of scales and ways of placement on adventitious bud induction

Table-1 shows that the regeneration ability of the outer layer scales of bulbil was better than the middle ones whose regeneration ability was better than the inner ones. The induction rate of the outer and the middle layer scales was higher (90.2 and 94.7%, respectively). The number of adventitious buds was more (9.4~11.1). By comparison, the induction rate of the inner layer was lower and the number of adventitious buds was less (715 and 2.3%, respectively). Besides, there were differences in contamination rate among three parts of scales. The sequence was the outer layer greater than the middle layer greyer than the inner layer.

That the adaxial side of Royal lily bulbils in contact with medium upward produced a higher callus induction rate, differentiation rate and larger number of adventitious buds than the adaxial side downward as shown in Table-2. Therefore, under the same conditions, the layment of adaxial side upward was the optimum layment for adventitious bud induction and adventitious bud growth in bulbil scale culture in vitro in Royal lily.

Effects of types and concentrations of cytokinin on adventitious bud induction:

Three types of cytokinin (BA, KT and ZT) can induce the outer and middle layer scales of bulbils to form adventitious buds. From the callus induction rate and differentiation rate, it can be seen that KT produced the highest callus rate and differentiation rate, BA produced the lowest and ZT was in the middle. In terms of the number of adventitious buds, KT was better than BA, which was better than ZT. Higher concentration of BA and KT were better for adventitious bud induction. But the lower concentration of ZT enhanced adventitious bud induction.

Especially, the explants cultured on MS medium supplemented with 1.0 mg L⁻¹ KT, 0.2 mg L⁻¹ NAA and 500 mg L⁻¹ CH exhibited the highest differentiation rate (100%), the largest number of adventitious buds (12.7) and the most sturdy adventitious buds as shown in Table-3.

By comparing differentiation rate and

growth status of adventitious buds, it could be concluded that the suitable medium for adventitious bud induction of Royal lily bulbils scales were MS basal medium added 1.0 mg L⁻¹ KT, 0.2 mg L⁻¹ NAA and 500 mg L⁻¹ CH. The results showed that inductive effect of inner layer scales of bulbils of Royal lily were worse than the outer and middle ones (Table-3 and Table-4). From the differentiation rate and number of adventitious buds, it can be seen that 1.5 mg L⁻¹ KT and 1.5 mg L⁻¹ ZT exhibited the higher differentiation rate (80~81.8%) with larger number of adventitious buds per explants (6~8.6). Although, 1.0 mg L⁻¹ BA exhibited the highest differentiation rate (92.3%), the number of adventitious buds was lowest (only 1.0) the appropriate medium for adventitious bud induction of inner bulbil scales from Royal lily was MS basal medium supplemented with 1.5 mg

L⁻¹ ZT and 500 mg L⁻¹ CH or MS basal medium supplemented with 1.5 mg L⁻¹ KT and 500 mg L⁻¹ CH.

Effects of BA, KT and NAA on multiplication in the multiplication cultures

Effects of different combinations of BA, KT and NAA on multiplication are shown in Table-5. The multiplication effects of different concentration combinations of KT and NAA were better than the combinations of BA and NAA. The multiplication coefficient (4.1~7.3) of the former was higher than the latter (1.22~3.4) and clumpy buds were sturdy. The medium supplemented with 0.5 mg L⁻¹ KT, 0.1 mg L⁻¹ NAA and 500 mg L⁻¹ CH exhibited the largest multiplication coefficient (7.3), the highest incidence rate of clump buds (74.24%) and the clumpy buds grew strongly. It could be concluded that MS supplemented with

Table 1. Comparison of different parts of scales of bulbils in adventitious bud induction

Parts of scales	No. of explants	Contamination rate (%)	Differentiation rate (%)	No. of adventitious buds (cm)
The outer layer scales	78	34.4	94.7	11.1
The middle layer scales	61	15.4	90.2	9.4
The inner layer scales	76	5.1	71.5	2.3

Table 2. Effects of different ways of placing on adventitious buds induction

Placing way	No. of explants	Calluses induction rate (%)	No. of calluses	Differentiation rate (%)	No. of adventitious buds (cm)	Length of adventitious buds (cm)
Adaxial side downward	56	71.9	7.9	71.9	8.2	0.4-2.0
Adaxial side upward	43	97.7	12.6	97.7	9.1	0.4-2.5

Table 3. Effects of types and concentrations of cytokinin on adventitious bud induction of outer and middle layer scales of bulbils from Royal lily

BA (mg L ⁻¹)	KT (mg L ⁻¹)	ZT (mg L ⁻¹)	No. of explants	Calluses induction rate (%)	No. of calluses	Differentiation rate (%)	No. of adventitious buds (cm)	Length of adventitious buds (cm)
1.0	0.0	0.0	66	84.20	9.10	86.40	8.00	0.7-1.2
1.5	0.0	0.0	62	87.50	14.8	87.50	12.7	1.5-1.7
0.0	1.0	0.0	68	100.0	14.6	100.0	12.7	2.5-2.6
0.0	1.5	0.0	63	100.0	9.90	100.0	10.9	0.4-0.8
0.0	0.0	1.0	64	95.80	10.3	95.80	11.3	0.8-2.0
0.0	0.0	1.5	66	90.80	5.80	90.80	10.2	0.2-1.2

0.5 mg L⁻¹ KT, 0.1 mg L⁻¹ NAA and 500 mg L⁻¹ CH was the optimum medium for multiplication culture of bulbil scales of Royal lily.

In addition, it was found that all the clumpy buds could develop 3~5 roots ranging 3~4 cm in length. The plantlets could be successfully transplanted after acclimatization without a rooting culture.

Effects of different parts of bulbil scales and ways of placement on adventitious bud induction

The sequence of regeneration ability of bulb was the outer layer scales greater than the middle layer scales greater than the inner layer scales¹⁶, which was similar to the results obtained in this experiment. The time of greening of the outer and the middle layer scales was almost in the next 6~10 days. Besides, the outer and the middle layer scales developed more adventitious buds and a larger number of calluses. Therefore, the outer and the middle layer scales are better explants in bulbil scale culture *in vitro* in Royal lily.

Affected by plant polarity, differentiation of bulbil scales was influenced by the way of placement on medium. The result showed that the adaxial side in contact with medium upward could induce more and stronger adventitious buds than the downward, which was in accord with the result gained in¹¹. The different regeneration ability of different parts of bulbil scales may be caused by different content of endogenous hormones and associated with different relative contents of growth promoting substances and growth inhibitors and controlled by balance of the two kinds of endogenous hormones⁴. It was speculated that there may be different contents of endogenous hormones in different parts of bulbil scales. It also, may have something to do with characteristics of physiology and biochemistry or different nutrient substances in bulbil scales.

Effects of plant growth regulators on induction and multiplication of bulbil scales of Royal lily

In previous reports of plant tissue culture

Table 4. Effects of types and concentrations of cytokinin on adventitious bud induction of the inner layer scales of bulbils from Royal lily

BA (mg L ⁻¹)	KT (mg L ⁻¹)	ZT (mg L ⁻¹)	No. of explants	Calluses induction rate (%)	Differentiation rate (%)	No. of adventitious buds (cm)	Length of adventitious buds (cm)
1.0	0.0	0.0	28	0.0	50.5	1.0	0.4-0.5
1.5	0.0	0.0	27	10.3	63.0	1.0	0.4-0.5
0.0	1.0	0.0	26	12.5	92.3	1.0	0.5-0.6
0.0	1.5	0.0	22	31.8	81.8	6.0	0.3-0.4
0.0	0.0	1.0	24	12.5	66.7	1.0	2.0-3.0
0.0	0.0	1.5	24	26.7	80.0	8.6	0.3-0.6

Table 5. Effects of BA, KT and NAA on multiplication in the multiplication culture

BA (mg L ⁻¹)	KT (mg L ⁻¹)	NAA (mg L ⁻¹)	No. of explants	The incidence rate (%) of clumpy buds	Multiplication coefficient (%)	Length of adventitious buds (cm)
0.5	0.0	0.1	62	41.9	1.6	3.0-6.0
0.5	0.0	0.2	63	57.1	1.9	2.5-6.0
1.0	0.0	0.1	92	15.2	1.2	3.0-6.0
1.0	0.0	0.2	71	39.4	1.7	2.0-4.0
2.0	0.0	0.1	71	71.7	3.4	2.0-4.0
2.0	0.0	0.2	79	53.2	2.7	2.0-5.0
0.0	0.5	0.1	66	74.2	7.3	4.0-6.05
0.0	0.5	0.2	78	70.5	4.4	4.0-5.5
0.0	1.0	0.1	57	66.7	4.1	3.0-6.0
0.0	1.0	0.2	51	76.5	5.4	5.0-6.0

on lily, a combination BA and NAA was widely used for shoot induction^{7,10,15} and proliferation^{2,3}. There was no report on the comparison of induction effect of BA, KT and ZT on plant tissue culture in *Lilium*. Studies were made of the effects of BA, KT and ZT on adventitious bud induction of bulbil from Royal lily in this experiment. The results showed that for the outer and middle layer scales, the higher BA concentration could induce adventitious buds efficiently.

But the lower concentration KT and ZT had a significant promotion on bulbil scale induction and growth, otherwise for the inner layer scales. BA had no significant promotion on bulbil scale's induction and growth and the higher KT and ZT concentration was better for adventitious bud induction. Therefore, it would be advisable to choose the suitable kinds of cytokinin and concentration according to material parts on culture in vitro of lily. In particular, it was the first time to study effects of ZT on adventitious bud induction in the present study.

Compared with the previous researches in which KT was used in combination with NAA for shoot proliferation of Royal lily's bulbil scales. More strong plantlets with higher multiplication coefficient than combination of BA and NAA were obtained in the present experiment.

It was found that the clump buds from all treatments demonstrated etiolation in different degree and finally died (death rate ranging from 3.22-50.77%), when subculture time was over 60 days, especially in the treatment with higher multiplication coefficient. The main reason may be that the subculture time was so long that the medium can't meet clump buds nutrient requirement. In addition, the more clump buds the greater respiratory intensity and the more metabolic substances produced by respiration. The experiment showed that the optimum subculture time of bulbil scales of Royal lily was 45 days.

It was also found that all of the shoots cultured on the proliferation media developed a number of sturdy and good quality roots. These rooted plants could be transplanted after acclimatization without rooting culture, which would be of a practical significant importance in promoting industrial seeding of Royal lily.

CONCLUSION

Bulbils from Royal lily were used as explants. A tissue culture and fast propagation system was established in this experiment, which would provide theoretical and technical basis for fast propagation, detoxification utilization and resource conservation of Royal lily and also, would be of a practical significant importance in promoting industrial seeding of Royal lily.

ACKNOWLEDGEMENTS

This research was partially supported by Payame Noor University of Iran and Islamic Azad University of Iran. We thank our colleagues from biotechnology laboratory that provided insight and expertise that greatly assisted the research, although they may not agree with all of the interpretations of this paper.

REFERENCES

1. Bacchetta, L., P.O. Remotti, C. Bernardini and Saccardof. Adventitious shoot regeneration from leaf explants and stem nodes of *Lilium*. *Plant Cell Tissue and Organ Culture*, 2003; **74**: 37-44. DOI: 10.1023/A:1023315321931.
2. Ding, L., GA. Liu, WD. Tian, XF. ling and Y.H. Zhang. Study on tissue culture and clonal propagation of *Lilium longiflorum* X *L. formosanum*. *J. Northwest Normal Univ.*, 2001; **37** (1).
3. Guo, HE. and M. Lei. The bulblet scale and bulblet culture in vitro of *Lilium lancifolium* Thumb. *Chinese Agric. Sci. Bull.*, 2006; **2**: 72-74.
4. Jin, S.M., LP. Yang, P. Lv and ER. Zhou. Changes of endogenous hormones in *Lilium pumilum*. 1. *Northeast For. Univ.*, 2005; **33** (1): 20-22.
5. Kapoor, R., S. Kumar and J.K Kanwar. Bulbil regeneration from we vitro root explant in lily hybrids. *Horticultural Sci*, 2008; **35**(3): 107-112.
6. Kumar, S., V. Awasthi and J.K Kanwar. Influence of growth regulators and nitrogenous compounds on in vitro bulblet formation and growth in oriental lily. *Horticultural Sci*, 2007; **34**: 77-83.
7. Kumar, S., J.K Kanwar and DR. Sharma. In vitro propagation of *Lilium*. *Adv. Horticultural Sci*, 2006; **20**: 181-188.
8. Kumar, S., M. Kashyap and DR. Shanna. In vitro regeneration and bulblet growth from lily bulb scale explants as affected by retardants,

- sucrose and irradiance. *Biol. Plantarum*, 2005; **49**: 629-632. DOI: 10.1007/s10535-005-0064-z.
9. Li, C.H., Z. Wu, Y. Yang and Y]. Long. Study on inducing and proliferating conditions of adventitious buds from bulblets of Yixing lily. *Acta Agric. ShangHai*, 2008; **24** (1): 27-31.
 10. Li, B.H., XL. J in and X.M. Liu. Establishment of regeneration system for bulbs of *Lilium* sp. *Jiangsu Agric. Sci*, 2008; **4**: 83-85.
 11. Li, H.Y., H.X. Ma, 1]. Lei, GH. Yu and J.T. Zhou. Study on fast propagation of oriental Lily from bulbet scale combined with blade segment. *Jiangsu Agric. Sci*, 2006; **3**: 91 -94.
 12. Lian, M.L., D. Chakrabarty and KY. Paek. Bulblet formation from bulb scale segments of *Lilium* using bioreactor system. *Biol. Plantarum*, 2003; **46**: 199-202. DOI: 1 0.1 023/A:1022890208500.
 13. Qiu, L.H., L.H. Luo, Q. Wang and L. Han. Tissue culture and fast propagation of *Lilium sulphureum* baker. *Chinese Wild Plant Resour*; 2004; **23** (2): 64-65.
 14. Sun, X.M., Q. Fu, P. Cao, HG. Yang and G.X. Ren. Studies on tissue culture of Royal lily . *J. Liaoning For. Sci. Technol*, 2001; **5**: 8-10.
 15. Tanaka, A., Y. Hoshi, K. Konodo and K. Taniguchi. Induction and fast propagation of shoot primordial from shoot apices of *Lilium japonicum*. *Plant Tissue Culture Letters*, 1991; **8**(3): 206-208.
 16. Wang, G., J. Du, G.Y. Li, W.F. Liang and HT. Xin. Study on tissue culture and clonal propagation of *Lilium davz'dz'z'* var. Unicolor (H 0og) Cotton and *Lilium browniz'* FEBrown ex Miellez. 1. *Northwest Normal Univ.*, 2002; **38**(1): 69-71.
 17. Xu, L.F., F.W. Ma and D. Liang,. Plant regeneration from in vitro cultured leaves of Lanzhou lily (*Lilium davz'dz'z'* var. unicolor). *Scientia Horticulture*, 2009; **119**: 458-461. DOI: 10.1016/j.scienta.2008.08.026.
 18. Xin, P.Y. and SB. Luo. Tissue culture and rapid propagation of *Lilium regale*. *Subtropical Plant Sci*, 2007; **4**: 57.
 19. Yao, Q.J., B. Xia, F. Peng, S.L. He and K. Zhou. Study on polyphenol oxidase isozyme of Royal lily and *Lilium sargentz'ae* in Minjiang River Valley. *Guihaia*, 2006; **26**(1): 92-96.
 20. Zhao, QF. and PX. Ma. Advances in detoxification of lily and identification of virus research. *Northern Horticulture*, 2007; **3**: 77-79.
 21. Zhang, S.J., F.L. Wang, HG. Zhou, R.L. Jiang. Tissue culture for fast propagation of Stargaz. *Jiangsu Agric. Sci*, 2004; **4**: 72-73.