

Laboratory Outbreak investigation of Sudden Death Syndrome in Broiler Chicken in Kathmandu Valley Nepal 2010

Kedar Karki

Senior Veterinary Officer, Central Veterinary Laboratory, Tripureswor Kathmandu (Nepal).

(Received: 25 October 2011; accepted: 02 December 2011)

The incidence of death of broiler birds above 40 days suddenly increased in the month of July-October 2010 in Kathmandu valley. Birds that were presented for post-mortem examination at the Central Veterinary Laboratory Tripureswor Kathmandu were usually found dead on their backs with wings out-stretched. Gross abnormalities recorded on post mortem examinations were: muscle oedema, pulmonary, renal and liver congestion, dark black to pale yellow streaked liver bile filled gall bladder and congestive splenomegaly, blood clot in atrium haemorrhage in duodenal muscle, whitish yellow pasty fluid in proventriculus gland, greenish coloration marked intact feed particles in gizzard and excessive mucous filled swollen intestine. Incidence rate was recorded between 1.5 to 2.5% of the flock. The mean mortality rate due to sudden death syndrome was 1.3 - 9.6%. *Penicillium* spp., *Aspergillus* spp., with CFU ranging from 56×10^4 - 62×10^5 , to uncountable mold count. *E. coli*, *Streptococcus* spp. and *Staphylococcus* spp. were the usual organisms isolated from culture samples of liver, lung, spleen and proventriculus. The condition seems to be related to mycotoxicosis. Reduction of mortality was achieved by feed restriction, with 8-10% reduction in nutrient density. Supplementation of glucose containing electrolyte, liquid toxin binder, Immunomodulator, acidifier and antibiotic therapy.

Key word: Sudden death syndrome, broiler birds, Kathmandu Valley, *Penicillium*, CFU, Ranging from 56×10^4 - 62×10^5 /gm, Uncountable mold count.

SDS has been recognized for over 30 years, and is also referred to as acute death syndrome or "flip-overs". It is most common in males when their growth rate is maximized. Mortality may start as early as 3-4 days, but most often peaks at around 3-4 weeks of age, with affected birds being found dead on their back. Mortality may be found at 1.5 to 2.0% in mixed-sex flocks and as high as 4% in male flocks only (George Qiao 2007) Ph.D. Poultry Nutritionist (www.championfeeds.com). Sudden death

syndrome has developed into a major problem to the broiler industry in many parts of the world. Broilers of all ages are affected starting as early as 2 days of age and continuing through to market age. Peak mortality usually occurs between 3 and 4 weeks of age. (Gardiner *et al.*, 1988). The syndrome has been reported to cause between 1.31 and 2.46 per cent mortality with males more commonly affected than females (Riddell and Orr, 1980), however Proudfoot and Hulan (1982) reported 0.90 to 3.61 per cent mortality due to SDS in broilers. From 0.71 to 4.07 per cent as reported by Riddell and Springer, (1985) where as Ononiwu *et al.*, (1979) reported 1.0 per cent mortality due to SDS in broilers. Brigden and Riddell (1975) reported that 70- 80 per cent of male mortality and 20-25 per cent of female broilers chickens mortality could be

* To whom all correspondence should be addressed.

attributed to acute death syndrome or "Flip-Over Disease". Thus, SDS is a leading contributor to mortality in broiler chicken production. Although the precipitating event has yet to be ascertained, "Cardiovascular failure" appears to be the immediate cause of death (Siddiqui, *et al.*, November 2009).

Background of outbreak of Sudden Death Syndrome in Kathmandu Valley

From the first week of July to October 2010 there was a sudden increase in mortality of broilers above 6 week age (Table 1) in Kathmandu valley. There were no premonitory signs. Just before death, birds appear normal and it is common to observe the birds use to eat, drink or walk normally. Then birds use to exhibit clinical signs such as extending their neck, squawk and start wing beating as well as leg extension before falling on their back.

Postmortem finding of SDS birds

Gross abnormalities recorded on post mortem examinations were muscle oedema, pulmonary, renal and liver congestion, dark black to pale yellow streaked liver bile filled gall bladder and congestive splenomegaly, blood clot in atrium haemorrhage in duodenal muscle, whitish yellow pasty fluid in proventriculus gland, greenish coloration marked intact feed particles in gizzard and swollen intestine with excessive mucous filled. All these post mortem observations conform to the descriptions of the syndrome made by Ononiwu *et al.*, (1979).

Laboratory finding of Mycobiota and Microbiota of Postmortem tissue samples

A total 176 tissue samples of lung, liver, spleen, proventriculus and gizzard, were collected during postmortem examination and were subjected for both bacterial and mycological culture. Results of microbiological examination are given in Table 2.

Treatment and Preventive measure given to the rest of birds in flock

All birds remaining in flocks were subjected to restricted feed up to 8-10% these percentages differ from the ones in the abstract, and fed twice daily. Supplementation with glucose containing electrolyte, liquid toxin binders, immunomodulator, and simple broad-spectrum antibiotics and acidifiers were provided in water. Vitamin B complex supplementation was totally withdrawn. All birds remaining in all affected farms responded well to the above management and there was a marked improvement in the overall condition of the flock.

RESULTS AND DISCUSSION

Sudden Death Syndrome (SDS) is an acute heart failure disease that affects mainly fast growing male chickens that seem to be in good condition. Although a common feature in fast growing birds, the pathogenesis remains unclear (Ononiwu *et al.* 1979). Cardiac arrhythmias are involved in the pathogenesis of SDS with ventricular arrhythmias (VA) being the most common observation representing premature ventricular contractions and fibrillation (Olkowski and Classen, 1997; 1998). It has been reported that broilers fed with high vitamin D3 diet above the recommended levels in an attempt to prevent commonly occurring leg problems were 2.5 fold more likely to succumb to acute heart failure and die of SDS (Nain *et al.* 2007). SDS was also experimentally induced by feeding diets containing the mycotoxin moniliformin that resulted into cardiac injury with subsequent alterations in cardiac electrical conductance (Reams *et al.*, 1997) suggesting the possible role of chronic mycotoxicosis to the causation of SDS. Other

Table 1. Epidemiology of Affected flock with sudden Death Syndrome in July-October 2010

Observation /Duration	no. of farmers/flock	Population at risk	Morbidity (%)	Mortality (%)	No. of samples examined
July	63	16620	4250 (25.57%)	369(2.22%)	63
August	51	15450	1235(7.99%)	232(1.50%)	51
September	32	10260	848(8.26%)	157(1.53%)	32
October	30	15700	2380(15.16%)	149(0.94%)	30
Total	176	58030	8713(15.01%)	907(1.56%)	176

Table 2.

No. of samples	Bacterial isolated	Fungi isolated	Positive no.	negative no.
176	<i>E.coli</i> , <i>Streptococcus</i> , <i>Staphylococcus</i>		35	141
176		<i>Aspergillus</i> , <i>Penicillium</i> ,	145	31

Rapid test for AI, ND, IBD

	No of samples	Positive	Negative
AI	20	0	20
ND	20	0	20
IBD	20	0	20

implicated causes of SDS include continuous artificial lighting (Ononiwu et al., 1979b), deviations in dietary calcium and phosphorus (Scheideler et al, 1995), feeding crumble-pellet diets (Proudfoot et al, 1982), dietary fat content (Rotter et al, 1985) and feeding frequency (Bowes et al, 1988). The latter recommendation of restricted feeding supports well the previous observation that abdominal fat deposition increases the risk of SDS such that restrictions on calorie:protein ratio decreases the incidence of SDS (Mollison et al., 1984).

Recommendation

The present investigation indicates that broilers in good body weight condition when not harvested timely and remaining in poultry shades for prolonged periods suffer stressful events and even die sudden. Also, it is possible that increased humidity and hot seasons favors the growth of mold and fungus in stored feeds increasing the risk of birds to mycotoxicosis. Detail Histopathological examination of affected organ need to be carried out for further verification of the involvement of mycotoxin for this syndrome.

REFERENCES

1. Bowes VA, R.J. Julian, S. Leeson and T. Stirtzinger., Effect of feed restriction on feed efficiency and incidence of sudden death syndrome in broiler chickens. *Poultry Science* **67**(7): 1102-4 (1988).
2. George Qiao: Reduction of Early Mortality in Broiler Chickens through Nutrition and Management: Champion Feed service limited: www.championfeeds.com pdf pp 1-2 (2007).
3. Mollison B., W. Guenter, and B.R. Boycott., Abdominal fat deposition and sudden death syndrome in broilers: the effects of restricted intake, early life caloric (fat) restriction, and calorie: protein ratio. *Poultry Science* **63**(6): 1190-200 (1984).
4. Nain S, B. Laarveld B, C. Wojnarowicz C, and A.A. Olkowski., Excessive dietary vitamin D supplementation as a risk factor for sudden death syndrome in fast growing commercial broilers. *Comparative biochemistry and physiology. Part A, Molecular & Integrative Physiology* **148**(4): 828-33 (2007).
5. Olkowski, A.A., C. Wojnarowicz, S. Nain, B. Ling, J. M. Alcorn, and B. Laarveld., A study on pathogenesis of sudden death syndrome in broiler chickens. *Research in Veterinary Science* **85**(1): 131-40 (2008).
6. Ononiwu, J.C., R.G. Thomson, H.C. Carlson, and R.J. Julian., Pathological Studies of "Sudden Death Syndrome" in Broiler Chickens. *Canadian Veterinary Journal* **20**(3): 70-73 (1979).
7. Ononiwu, J.C., R.G. Thomson, H.C. Carlson, and R.J. Julian., Studies on effect of lighting on "Sudden death syndrome" in broiler chickens. . *Canadian Veterinary Journal* **20**(3):74-7 (1979b).
8. Olkowski A.A. and H.L. Classen., High incidence of cardiac arrhythmias in broiler chickens. *Zentralblatt für Veterinärmedizin. Reihe A* **45**(2):83-91 (1998).
9. Olkowski A.A. and H.L. Classen., Malignant ventricular dysrhythmia in broiler chickens

- dying of sudden death syndrome. *Veterinary Record*. **15**: 140(7): 177-9 (1997).
10. Proudfoot FG, H.W. Hulan, K.B. McRae., The effect of crumbled and pelleted feed on the incidence of sudden death syndrome among male chicken broilers. *Poultry Science*. **61**(8): 1766-8 (1982).
 11. Reams R.Y, H.L. Thacker, D.D. Harrington, M.N. Novilla, G.E. Rottinghaus, G.A. Bennett , and J. Horn., A sudden death syndrome induced in poults and chicks fed diets containing *Fusarium fujikuroi* with known concentrations of moniliformin. *Avian Disease*. **41**(1):20-35 (1997).
 12. Rotter B, W. Guenter, and B.R. Boycott., Sudden death syndrome in broilers: dietary fat supplementation and its effect on tissue composition. *Poultry Science*. **64**(6): 1128-36 (1985).
 13. Scheideler SE, D.V. Rives, J.D. Garlich, and P.R. Ferket., Dietary calcium and phosphorus effects on broiler performance and the incidence of sudden death syndrome mortality. *Poultry Science* **74**(12): 2011-8 (1995).
 14. Siddiqui, M. F. M. F.*, M.S. Patil, K.M.Khan, and L.A.Khan : Sudden Death Syndrome – An Overview: *Veterinary World*, **2**(11): 444-447: (2009).