

Prevalence and Salient Morphological Features of Myiasis Causing Dipteran Flies in Jeddah, Saudi Arabia

Masroor Alikhan^{1*}, K. Al Ghamdi², Fahad S Al Zahrani¹,
Emad I Khater^{1,3} and Ahmad M Allam^{1,4}

¹Public Health Pest Laboratory, Al Amana, Bariman, Jeddah, KSA.

²Department of Biological Sciences, K. Abdul-Aziz University, Jeddah, KSA.

³Department of Entomology, Faculty of Science, Ain Shams University, Cairo, Egypt.

⁴Department of Parasitology and Animal Diseases, Veterinary Research Division, National Research Centre, Egypt.

<http://dx.doi.org/10.13005/bbra/2612>

(Received: 06 February 2018; accepted: 23 February 2018)

A number of dipteran fly species are found in Jeddah with public health and veterinary importance due to their association with myiasis of livestock and humans. Although several reports are available on the myiasis causing flies from different parts of Saudi Arabia very little is known available on prevalence and characterization of these flies from Jeddah Province, which include the major harbour on the Red Sea, western Saudi Arabia and acts the gateway for millions of live animals in the kingdom. Therefore to fill this knowledge gap, the objective of this work is to fill up this gap. For fly survey, the adult flies were collected by the means of Malaise & Bait traps and sweeping nets from different marked locations including slaughter houses and live animals farms. The total number of flies from each locality was counted and were sorted for taxonomic species identification using specific pictorial keys. In this study, ten dipteran fly species belonging to six families were identified as causing myiasis on the basis of reported cases in Saudi Arabia. The flies collected during the survey were *Megaselia scellaris*, *Musca domestica*, *Fanniia canicularis*, *Sarcophaga haemorrhoidalis* (*Bercaea cruentata*); *Parasarcophaga ruficornis*, *Wohlfahrtia nuba*, *Chrysomya marginalis* *C. albicep*, *C. megacephala*, and *Oesteris ovis*. Among collected flies the highest number were of *Musca domestica* (67.6%) and the rest were other species. Most of the species were collected from slaughter houses (50%) followed by animal farms (39%). The present study identified a number species of myiasis-causing flies of the public health and veterinary significance, which should inform wider studies on their seasonal abundance, significance and consequent implementation of preventive control measures.

Keywords: Diptera, Flies, Myiasis, Morphology, Prevalence.

Myiasis is defined as parasitic infestation of the body organs of vertebrate animals and humans by a variety of fly species maggots that feed upon living or dead tissues (Zumpt, 1965)

Myiasis can be classified as anatomical which includes open wounds, furuncular, intestinal

or cavity myiasis or ecological which includes obligatory, facultative and accidental (Pseudo myiasis) (Derraik et al 2010).

Obligatory or specific myiasis is caused by parasitic flies which require living tissues for their larval development. Facultative or semi-

*Corresponding author E-mail: alikhansmasroor@hotmail.com

This is an Open Access article licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License (<https://creativecommons.org/licenses/by-nc-sa/4.0/>), which permits unrestricted Non Commercial use, distribution and reproduction in any medium, provided the original work is properly cited.



specific myiasis is caused by opportunistic flies which live freely and lay eggs on the decaying organic matter and invade the living tissues especially pre-existing wounds. Accidental or pseudo-myiasis occurs when the food contaminated with fly eggs or larvae is ingested by host animals which subsequently develop pathological reactions (Hall, 1991).

Myiasis is distributed worldwide especially in poor socioeconomic regions of tropical and subtropical countries (Francesconi, and Lupi, 2012). From Saudi Arabia many workers reported different forms myiasis in vertebrate animals and humans (Omar & Abdullah, 1992; Badawi 1994; Fatani & Hilali 1994; Hall & Wall 1995; Khayat 2002; Al Ahmad et al 2006; Wakid 2008; Bosly 2013 and Zagloul et al 2013,) however, there is a paucity of literature on the identification and the habitats of myiasis causing flies found in Jeddah which is a large commercial city and harbour, through which where millions of livestock enter the kingdom annually and a metropolis with large number of animal sheds. Therefore, the objective of this study was to identify the myiasis causing flies in Jeddah and characterize the habitats. The results will provide important information that will enable the veterinary and public health authorities to identify the flies rapidly and accurately and to take appropriate actions for myiasis prevention and control.

MATERIAL AND METHODS

Flies were collected from slaughter houses (abattoirs), animal farms, fish and vegetable markets of Jeddah city. Jeddah is a metropolis and a commercial city and harbour located on the west coast of Red Sea. (Latitude 21.29 N and Longitude 39.7 E). The climate is usually moderate except during summer when the temperature is high

which might reach 45°C, and winter is moderately temperate. The average high is approximately 35°C and low average is about 23°C. The average relative humidity is approximately between 50% to 60%.

myiasis causing adult flies were collected by the means of Malaise traps, bait traps and sweeping nets from previously marked locations. These locations included slaughter houses located in the districts of municipalities: Abhur, Al Mattar, Um Elsalam, Al Balad and Al-Janoub (Table-1), while the animal farms especially sheep, goats, camel & horse paddocks located in South Jeddah, East of Harmain Road, Al Nuzah, Um Elsalam, Al Jamia, Asfan and Dhahban districts.

After preservation adult flies were morphologically characterized and classified with the help of available authentic keys. (Akbarzadeh et al 2015; Al ahmadi and Salem et al 1999, Al Ghamdi and Alikhan 2015; Alikhan et al 2016.; Dabbour and El Dawy 1981, Irish et al. 2014; Mc Alpine 1989; Nazni et al 2011; Pont 1991; Setyaningrum and Al Dhafer, 2014; Szpila 2012; Verves 2005).

In the present paper we mentioned only salient morphological features of the flies. The detailed identification key was already published in previous paper (Alikhan et al 2016).

RESULTS AND DISCUSSION

According to morphological identification of adult flies the following ten dipteran fly species belonging to six families were recorded as myiasis causing on the basis of reported cases in Saudi Arabia (Table.3). The flies collected during survey were *Megaselia scalaris*, *Musca domestica*, *Fanniia canicularis*, *Sarcophaga haemorrhoidalis* (*Bercaea cruentata*); *Parasarcophaga ruficornis*, *Wohlfahrtia nuba*, *Chrysomya marginalis*, *chrysomya albiceps*, *Chrysomya megacephala*,

Table 1. Showing the approved Abattoirs in Jeddah

Municipality	District	Geographic coordinates	
UmElsalam	Al-nakheel	39.25935	21.52699
Al- janoub	Al- khumra	39.22224	21.38080
Abhor	Al- nahda 1	39.13383	21.60995
Al- balad	Al- sabeel 1	39.20245	21.48317
Al -matar	Al- nozha 3	39.18270	21.62562

Table 2. Showing total number of flies and their percentage collected from different locations

Family Name	Species Name	Abattoirs Collection	Animal farm Collection	Fish & veg. Market Collection	Animal Carcasses Collection	Total No. Of flies	Total percentage
Phoridae	<i>Megasellia scalaris</i>	56	62	6	82	206	7.721%
Muscidae	<i>Musca domestica</i>	980	766	32	26	1804	67.616%
Fanniidae	<i>Fanniia canicularis</i>	18	11	6	0	35	1.311%
Sarcophagidae	<i>Sarcophaga haemorrhoidalis</i>	106	15	59	53	233	8.733%
Sarcophagidae	<i>Parasarcophaga ruficornis</i>	18	10	4	5	37	1.386%
Sarcophagidae	<i>Wohlfahrtia nuba</i>	14	5	8	4	31	1.161%
Calliphoridae	<i>Chrysomya albiceps</i>	120	49	79	28	276	10.344%
Calliphoridae	<i>Chrysomya megacephala</i>	6	3	2	1	12	0.449%
Calliphoridae	<i>Chrysomya marginalis</i>	13	6	4	3	26	0.974%
Oestridae	<i>Oestrus ovis</i>	3	5	0	0	8	0.299%
TOTAL		1334	932	200	202	2668	
Percentage		50%	34.93%	7.50%	7.57%		

and *Oesteris ovis*. The present prevalence of each fly species was recorded (Table.2). The highest prevalence was recorded for the common house fly *Musca domestica* (67.6%). Most of the species were collected from slaughter houses (50%) and animal farms (39%). (Table.2).

Following are the salient morphological diagnostic characteristics and habitats of the species collected.

Family: Phoridae

Megasellia scalaris (Loew, 1866) (Coffin fly or scuttle fly)

Brownish small fly which is also named as scuttle fly due to its sudden and rapid movements. The adult fly may be easily recognised by “humpbacked” thorax and laterally compressed hind femur. The fore margins of the wings are thickened, wing veins are parallel to each other and without cross veins. One segmented antenna with long arista.

The fly maggots are found near rotten meat, vegetable remains, trash containers, abattoirs, over animal carcasses and near poor sanitary conditions. (it causes human accidental myiasis in wounds, intestine, eyes, respiratory system and urino-genital system).




Fig.1. A map of Jeddah city showing different locations of Flies collection

Table 3. Cases of human and animal myiasis recorded from Saudi Arabia

IN HUMANS Family	Species	Type of Myiasis	Reported By	Region
Phoridae	<i>Megaselia scalaris</i>	Urinary	Wakid, MH 2008	Jeddah
Sarcophagidae	<i>Wohlfahrtia magnifica</i>	Aural/Scalp	Al Jbar, I 2015; Al Badri et al 2016	Al Ahsa
Sarcophagidae	<i>Sarcophaga</i> sps.	Open Diabetic wound	Zaglool et al 2013	Makkah
Calliphoridae	<i>Carylobia anthropophaga</i>	Cutaneous	Omar & Abdallah 1992; Afifi et al 2015	Al Asir/Al Baha
Oestridae	<i>Dermatobia hominis</i>	Cutaneous	Al Otaibi 2016; Akhtar et al 2000; Hiba Radwan 2015	Al Khobar/Al Asir/Taif
Oestridae	<i>Ostris ovis</i>	Ophthalmomyiasis	Kenway et al. 2014	Al Asir
IN ANIMALS Oestridae	<i>Ostris ovis</i>	Nasopharyngeal	Al Ahmad et al. 2000; Banja & Madbouly 1981; Bosly 2013	Riyadh/West/Jazan
Oestridae	<i>Cephalopina titillator</i>	Nasopharyngeal	Fatani & Hilali 1994; Ghandour, 1988; Banja & Madbouly 1981; Buttiker & Zumpt 1982	Al Ahsa/West/West/Riyadh
Calliphoridae	<i>Chrysomya bezziana</i>	Dermal	Al Ahmad et al 2001, 2006	Riyadh
Calliphoridae	<i>Chrysomya albiceps</i>	Dermal	Al Ahmad 2001	Riyadh
Calliphoridae	<i>Chrysomya megacephala</i>	Dermal	Ramdan & El Bihari 1980	Hafuf
Sarcophagidae	<i>Wohlfahrtia magnifica</i>	Dermal	Al Ahmad 2001	Riyadh

Family: Muscidae***Musca domestica* (Linnaeus, 1758) (House fly)**

Thorax with 4 black stripes on greyish background, abdomen with yellowish spots on the anterior lateral sides below the wings and a dark midline. Antennae are short and 3 segmented, anal cell in the wing is closed, 4th vein angled with discal cell.

It lives close to humans and contaminate the food, feed and breed in garbage and animal faeces. (It is an agent of accidental myiasis of intestine and urino-genital system).

Family: Fanniidae***Fannia canicularis* (Linnaeus, 1761) (Lesser house fly)**

The thorax is brown-grey with less distinct three black brown stripes, abdomen yellow in colour, eyes are with white border in male, head silvery in colour, halteres are yellowish, the 4th vein not angled but straight, anal vein short.

High number of *Fannia* found in poultry farms and animal sheds. Deposit eggs on decaying organic matter such as cow dung, human and poultry faeces.

(It causes accidental myiasis of intestine and urino-genital system)

Family: Sarcophagidae***Sarcophaga haemorrhoidalis* (Fallen, 1817) (red-tailed flesh fly)**

Thorax with 3 dark longitudinal stripes, antennal arista is plumose at the base and bare at the tip; antenna and palpi are greyish black with white hair on gennae. Abdomen with grey black checker board. Sixth tergite of female is red in

colour, separated with a row of strong bristles at the edge.

Flesh flies usually attracted to rotten meat, fish, vegetables, colonize the corpses, animal and human faeces.

Larviparous (Lay larvae) (Produce myiasis on necrotic or dead flesh in both humans and animals.).

***Parasarcophaga ruficornis* (Fabricius, 1794) (flesh fly species)**

Can be identified by white hairs on gennae below eyes, antenna and palpi orange in colour. Breed both in faeces and dead bodies.

(breeds on flesh causing myiasis or carrion and faeces, also a vector for diseases).

***Wohlfahrtia nuba* (Wiedemann, 1830) (spotted flesh fly)**

Greyish with three black round spots at the end of each abdominal segment, antennal arista bare or with very short hairs, larviparous. Breed on the open wounds of the camel and causes severe myiasis.

(Causes severe myiasis in Camel it is a secondary invader of wounds.)

Family: Calliphoridae***Chrysomya albiceps* (Wiedemann, 1819) (Blow fly species)**

Body metallic green with narrow dark lines on the rear edge of each abdominal segment; anterior thoracic spiracle white/pale yellow, antennal arista plumose, head with silvery hair.

(Female deposit egg on the decomposing tissues or in the wounds of living in animals and in



Fig. 2. Myiasis in a sheep



Fig. 3. Myiasis in Human

humans and causes a specific myiasis in wounds, sores and nasal cavities.)

***Chrysomya megacephala* (Fabricius,1794) (oriental latrine fly)**

Large red eyes with metallic blue/green body, buccal area and gennae with orange setae. Anterior thoracic spiracle black/brown. Lower

calypter white at the base and the rest is dark.

(Causes myiasis in animals and humans resulting into economic losses in cattle and fish industries all over the world.)

***Chrysomya marginalis* (Wiedemann,1830)**

Anterior wing margins dark, anterior spiracle white in colour. Greater ampulla with thick




Fig. 4. Percent variation in the population density of adult flies causing myiasis in Jeddah




Fig. 5. Percentage of flies causing myiasis collected from different locations of Jeddah

hairs dorsally. Lower calypter with dense hairs.

Family: Oestridae

***Oestrus ovis* (Linnaeus, 1758) (Sheep nasal bot fly)**

Short stout and dark grey body with black spots at the abdomen. Head and legs are dull yellow.

Female fly deposit eggs in the nostrils of the camel or sheep. Usually found where sheep camel or goats are reared.

(Responsible for myiasis of skin, mucous membranes of mouth, eyes or nasal cavities both in animals and humans.)

Most of the reports on myiasis in animals & humans from Saudi Arabia are about the case studies. (Afifi *et al.*, 2015; Al Otaibi *et al.* 2016; Al Badri, 2016; Al Ahmad *et al.*, 2006;). Many myiasis causing flies collected by different workers from other parts of the country such as, *Cordylobia anthropophaga*, *Dermatobia hominis*, *Wohlfahrtia magnifica*, *Cephalopina titillator*, *Chrysomya bezziana* *etc.*, may be present in Jeddah region but could not be trapped during this study. Some of the workers studied the identification of forensic important flies from Jeddah (Al Ghamdi and Alikhan, 2015, Al shareef H. 2016) but there is no record of myiasis causing flies from this region.

CONCLUSION

Present study is important because it provides a list of flies causing myiasis to animals and humans in this part of the kingdom and responsible for the economic losses and agony. The flies may be different taxonomically but they affect animals and humans in the same way and their ecology and habitats also similar.

The easy method of identification is needed for the veterinary & health workers to control the myiasis causing flies and to reduce the economic loss and human sufferings.

ACKNOWLEDGEMENT

The authors are indebted to the AlAmana Public Health Laboratory, Jeddah for providing the facilities to carry out this research work. We are also thankful to all our colleagues for their kind help and suggestions during this study.

REFERENCES

1. Afifi MA, Jiman-Fatani AA, Alsiny FI, Anshasi WS. A new focus of autochthonous transmission of *Cordylobia anthropophaga* in Saudi Arabia. *Journal of Microscopy and Ultrastructure.*; 2015; **3**(2):82-5
2. Akbarzadeh k; Wallman J; Sulakova H; SDzpila K. Species identification of Middle Eastern blow flies (Diptera: Calliphoridae) of forensic importance. *Prasitol Res*, 2015; **114**: 1463-1472
3. Akhter J, Qadri SM, Imam AM. Cutaneous myiasis due to *Dermatobia hominis* in Saudis. *Saudi medical journal*. 2000; **21**(7):689-91
4. Al Ahmad, AM; Al Dawood, AS; and kheir, SM. Seasonal activity of flies causing myiasis in livestock animals using sticky traps baited with swarm lure-4 in Riyadh region, Saudi Arabia, *Scientific Journal of King Faisal University (Basic and applied sciences)*, 2006; **7**:107-119.
5. Alahmed, A. M. Incidence of myiasis in sheep caused by *Chrysomya bezziana* in Saudi Arabia. *J. King Saud Univ.*, **14**: Agric. Sc 2001; (2), 109-112.
6. Al Ahmadi AZ and Salem MM. *Entomofauna of Saudi Arabia, Part 1: Check list of Insects*. Academic Publishing & Press. King Saud University 1999.
7. AlBadri R, Harbi AIT, Tonnsi A, Almatary A, Hassanein R. Cutaneous Myiasis in a Child Scalp Caused by *Wohlfahrtia magnifica* (Diptera: Sarcophagidae): A Case Report. *MOJ Clin Med Case Rep*, 2016; **4**(3): 00093. DOI: 10.15406/mojcr.2016.04.00093
8. Al Jabr, Ibrahim. Aural Myiasis, a Rare Cause of Earache; *Case Reports in Otolaryngology*, 2015; Article ID 219529, 3
9. Al-Shareef, Species Identification of Forensic Important Blowflies (Diptera, Calliphoridae) that occur in Jeddah, Kingdom of Saudi Arabia, *Glob J Nurs Forensic Stud*, 2016; 1:3.
10. Amoudi MA , Diab FM, Abou-Fannah SSM. The occurrence of *Megaselia scalaris* (Loew) (Diptera: Phoridae) in saudi Arabia with some aspects on the life history and distribution in Riyadh Province. *J King Saud Univ.Sci* 1989; **1-2**: 43-51.
11. Amoudi MA. New records of some Sarcophagid flies with distribution of all known flesh flies (Diptera: Sarcophagidae) of Saudi Arabia. *J. of Egyptian Society of Parasitology*, 1993; **23**(1): 297-304.
12. Alikhan *et al.* Public health and veterinary important flies (order: Diptera) prevalent in Jeddah Saudi Arabia with their dominant characteristics and identification key. *Saudi*

- Journal of Biological Sciences*, 2016: <http://dx.doi.org/10.1016/j.sjbs.2016.08>.
13. Al-Ghamdi ; Alikhan et al. Characterization of forensically important Nicrophorus Flies (Diptera) of Jeddah, *Saudi Arabia. Advances in Environmental Biology*, 2015; **9**(8); 58-71.
 14. Alotaibi S, Al shahrani D, Omar Alzomor. Cutaneous Myiasis in Saudi Infant: *A Rare Case Report IMJH*, 2016; **2**(9): 8-12
 15. Badawi A. Arthropods of Medical and veterinary importance in the kingdom of Saudi Arabia. King Saud Uni. Press Riyadh 1994.
 16. Areej O Bakhraibah. Prevalence and organ distribution of larval *Oestrus ovis* (Diptera : Oestridae), *Cysticercus tenuicollis* and *Echinococcus granulosus* (Cestoda : Taenidae) in slaughtered sheep and goats at Jeddah in Saudi Arabia 2016.
 17. Banja A & Madbouly MM. Field and laboratory observations on three dipterous larvae causing myiasis in the western region of Saudi Arabia. B ulletin Faculty of Science, King Abdulaziz University, 1981; **5**: 77-80
 18. Bosly A. Hanan. Seasonal prevalence of *Oestrus ovis* L. (Diptera:Oestridae) larvae in infested sheep in Jazan Region,Saudi Arabia; *J. Parasitol. Vector Biol.*, 2013; **5**(5): pp. 66-71,
 19. Boorman J. Insects of Saudi Arabia Culicoides (Diptera: Ceratopogonidae) of the Arabian peninsula with notes of their medical & veterinary importance, *Fauna of Saudi Arabia*. 1989; **10**: 160-225.
 20. Buttiker W & Zumpt F. Myiasis in domestic animals. *Fauna of Saudi Arabia*, 1982; **4**: 520-524
 21. Dabbour, A.I., "Short note on dipterous flies in western and central regions of Saudi Arabia", *Journal of Agriculture Research*, Riyadh University, 1979; **4**: 81-83.
 22. Dabbour, A.I., Note on dipterous flies in western and central regions of Saudi Arabia", *Journal of Faculty of Science, Riyadh University*, 1979; **10**: 117-119.
 23. Dabbour AI and El Dawy M. Morphological and classification studies of some dipterous flies in the kingdom of Saudi Arabia. *Agriculture Research Centre Bulletin* 1981.
 24. Dabbour, A.I., Hammad, S.M., Insect and Animal pests and their control in the Kingdom of Saudi Arabia. Uni. Libraries, King Saud Univ., Riyadh (In Arabic) 1982.
 25. Fabio Francesconi and Omar Lupi Myiasis, *Clinical Microbiology Reviews*, 2012; **5**(1)
 26. Fatani A, Hilali m., Prevalence and monthly variations of the second and third instars of *Cephalopina titillator* (Diptera: Oestridae) infesting camels (*Camelus dromedarius*) in the Eastern Province of Saudi Arabia 1994 . -
 27. José G B Derraik, Allen C G Heath, Marius Rademaker. Human myiasis in New Zealand: imported and indigenously-acquired cases; the species of concern and clinical aspects *NZMJ*, 2010; **123** No 1322; ISSN 1175 8716
 28. Ghandour, A. M. Health Hazards in Humans and Animal Caused by Imported Livestock Diseases in Saudi Arabia. *Fauna of Saudi Arabia*, 1988; **9**: 468- 476.
 29. Hall, MJR. Screw worm flies as agents of wound myiasis; World animal Review, Special issue" New World Screw worm; Response to an emergency" R.D.S. Branckaert (Ed), 1991; **52** Pp.
 30. Hall MJR and Wall R. Myiasis of humans and domestic animals. *Adv. Parasitol*, 1995; **35**:257–334
 31. Hiba M. Radwan. Myiasis. *Trop Med Surg*, 2015; **3**(4): 1000197
 32. Irish S, Lindsay T, Wyatt N. Key to adults of Afro-tropical species of the genus *Chrysomya* Robineau-Desvoidy (Diptera: Calliphoridae). *African Entomol*, 2014; **22**:297–306
 33. James MT. *The flies that cause myiasis in man*. United states department of agriculture: Miscellaneous publication No. 631. Washington, DC: US Government Printing Office 1947.
 34. Kamal AS. Comparative study of thirteen species of sarcophagus Calliphoridae and Sarcophagidae (Diptera): *Bionomics. Ann, Entomol .Soc. Am*, 1958; **51**: 261-271.
 35. Kenway, MA; Al ashry HA; Shobrak M. Synanthropic flies of Asir Province, Southwest of Saudi Arabia. *J. of Ento and Acro Res.*, 2014; **46**(3).
 36. Khyat RM. A case report on oral myiasis in Saudi Arabia. *Saudi.Dent.J*, 2002; **14**: 140-2
 37. Majed H. Wakid. A Laboratory-Based Study for First Documented Case of Urinary Myiasis Caused by Larvae of *Megaselia scalaris* (Diptera: Phoridae) in Saudi Arabia 2008.
 38. Mashaly A, A preliminary study of forensically important necrophagous Diptera in Riyadh, Saudi Arabia. Conf. 8th Int. Cong. Of Dipterology, Postdam, Gremany 2014.
 39. Mc Alpine JF. Phylogeny and classification of the Muscomorpha. In: Mc Alpine JF, Wood DM (eds,) Manual of Nearctic Diptera 3. *Research Branch, Agriculture Canada Monograph*, 1989; **32**: 1397-1518
 40. Morsy,TA,; Fayad ME; Salama,MMI; Sabry,AA; El Seroagi AOM and Abdullah, KF. Some myiasis producers in Cairo and Giza abattoirs, *J Egypt Soc Parasitol.*, 1991; **21**(2) 539-546

41. Nazni WA, Jeffery J, Heo CC, Chew WK, Lee HL. Illustrated keys to adult flies of forensic importance in Malaysia. Institute for Medical Research, Kuala Lumpur 2011.
42. Omar MS and Abdullah RE. Cutaneous myiasis caused by tumbu fly larvae, *Cordylobia anthropophaga* in South-western Saudi Arabia. *Trop. Med. Parasitol*, 1992; **43**: 128-9.
43. Patton WS. Some notes on Indian Calliphorinae III *Chrysomya megacephala* Fabr. *Bulletin of Entomological Research*, 1922; **13**(1): 109-113.
44. Patton WS and Cushing EC. Studies on the higher dipteral of medical and veterinary importance: a revision of the genera of subfamily Calliphorinae based on a comparative study of the male and female terminalia. *Tropical Medicine and Parasitology*, 1934; **28**:123-130.
45. Pont AC, A review of Faniidae and Muscidae of Arabian Peninsula. *Fauna of Saudi Arabia*, 1991; **12**: 312-365.
46. Ramadan, R. O. and El Bihari, S. Dermal myiasis in animal farm in El Hofuf Area. Fourth Symposium on the biological aspects of Saudi Arabia. Saudi Biological Society, Faculty of Science, University of Riyadh, 1980; March 10-13, 1980; p 85.
47. Setyaningrum H & Al Dhafer HM. The calliphoridae the blow flies (Diptera : Oestroidea) of Kingdom of Saudi Arabia. *Egypt Acad. J. Biolog. Sci.*, 2014; **7**(1)49-139.
48. Sundharam J, Al-Gamal M. Myiasis in Saudi Arabia. *Annals of Saudi medicine*. 1994; **14**(4): 352.
49. Szpila K. Key for identification of European and Mediterranean blowflies (Diptera, Calliphoridae) of medical and veterinary importance – adult flies. In: Gennard D (ed) *Forensic entomology, an introduction*, II edition. Willey-Blackwell, 2012; pp. 77–81 + plates 5.1–5.9.
50. Verves YG. A catalogue of Oriental Calliphoridae (Diptera). *Dipterological Res*, 2005; **16**: 223–310
51. Zagloul DA, Tayeb K, Khodari YA, Farooq MU. First case report of human myiasis with *Sarcophaga* species in Makkah city in the wound of a diabetic patient. *J Nat Sc Biol Med*; 2013; **4**: 225-8
52. Zumpt F, Myiasis in man and animals in the Old World: A textbook for physicians, veterinarians and zoologists. London: Butterworths 1965.